arXiv:1303.4035v1 [math.ST] 17 Mar 2013

On the sphericity test with large-dimensional observations

Qinwen Wang[*] **and Jianfeng Yao**[†]

Qinwen Wang
Department of Mathematics
Zhejiang University
e-mail: `wqw8813@gmail.com`

Jianfeng Yao
Department of Statistics and Actuarial Science
The University of Hong Kong
Pokfulam, Hong Kong
e-mail: `jeffyao@hku.hk`

Abstract: In this paper, we propose corrections to LRT and John's test for sphericity in large-dimension. New formula for the limiting parameters in the CLT for linear spectral statistics of sample covariance matrices with general fourth moments are established. Using these formula, we derive the asymptotic distribution of the two proposed test statistics under the null without assuming parametric distribution for the population. Besides, simulations are conducted to analyze finite-sample performance of the corrected tests.

AMS 2000 subject classifications: Primary 62H15; secondary 62H10.

Keywords and phrases: Large-dimensional data, Sphericity test, John's test, CLT for linear spectral statistics, Large-dimensional sample covariance matrix.

1. Introduction

Consider a sample $Y_1, \ldots, Y_n$ from a p-dimensional multivariate distribution with covariance matrix Σ_p. An important problem in multivariate analysis is

---

[*] Research of this author was partly supported by the grants NSFC-11071213, zjnsfc-r6090034 and dpme-j20110031.

[†] Research of this author was partly supported by a HKU start-up grant.

to test the sphericity of such a multivariate distribution, namely the hypothesis $H_0 : \Sigma_p = \sigma^2 I_p$ where σ^2 is *unspecified*. If the observations represent a multivariate error with p components, the null hypothesis expresses the fact that the error is cross-sectionally uncorrelated (independent if in addition they are normal) and have a same variance (homoscedasticity).

Much of the existing theory about this test has been exposed first in details in Mauchly, J.W. (1940) about Gaussian LRT and later in John (1971), John (1972), Sugiura, N. (1972) and also in textbooks like Muirhead (1982, Chapter 8) and Anderson (1984, Chapter 10). Assume for a moment that the sample has a normal distribution with covariance matrix Σ_p (as far as the covariance matrix Σ_p is concerned, we will assume in all the paper that the mean is zero). Let be the sample covariance matrix $S_n = n^{-1}\sum_i Y_i Y_i^*$ and denote its eigenvalues by $\{\ell_i\}_{1\le i\le p}$. Two well established procedures for testing the sphericity are the likelihood ratio test and a test devised in John (1971). The likelihood ratio statistic is, see e.g. Anderson (1984, §10.7.2),

$$L_n = \left(\frac{(\ell_1 \cdots \ell_p)^{1/p}}{\frac{1}{p}(\ell_1 + \cdots + \ell_p)} \right)^{\frac{1}{2}pn} ,$$

which is a power of the ratio of the geometric mean of the sample eigenvalues to the arithmetic mean. It is here noticed that in this formula it is necessary to assume that $p \le n$ to avoid null eigenvalues in (the numerator of) L_n. If we let $n \to \infty$ while keeping p fixed, classical asymptotic theory indicates that under the null hypothesis, $-2\log L_n \Longrightarrow \chi^2_f$, a chi-square distribution with degree of freedom $f = \frac{1}{2}p(p+1) - 1$. This asymptotic distribution is further refined by the following Box-Bartlett correction:

$$P(-2\rho \log L_n \le x) = P_f(x) + \omega_2 \left\{ P_{f+4}(x) - P_f(x) \right\} + O(n^{-3}) ,$$

where $P_k(x) = P(\chi^2_k \le x)$ and

$$\rho = 1 - \frac{2p^2 + p + 2}{6pn}, \qquad \omega_2 = \frac{(p+2)(p-1)(p-2)(2p^3 + 6p^2 + 3p + 2)}{288p^2n^2\rho^2} .$$

The test procedure based on this correction will be hereafter referred as the *LRT test.*

By observing that the asymptotic variance of $-2\log L_n$ is proportional to $\operatorname{tr}\{\Sigma(\operatorname{tr}\Sigma)^{-1} - p^{-1}I_p\}^2$, John (1971) proposed to use the statistic

$$T_2 = \frac{p^2 n}{2} \operatorname{tr}\left\{S_n(\operatorname{tr} S_n)^{-1} - p^{-1}I_p\right\}^2 .$$

It is observed that T_2 is proportional to the square of the coefficient of variation of the sample eigenvalues, namely

$$T_2 = \frac{np}{2} \cdot \frac{p^{-1}\sum(\ell_i - \overline{\ell})^2}{\overline{\ell}^2} , \qquad \text{with } \ \overline{\ell} = \frac{1}{n}\sum_i \ell_i .$$

Following the idea of the Box-Bartlett correction, Nagao (1973) established an expansion for the distribution function of the statistics T_2,

$$\begin{aligned} P(T_2 \le x) \quad &= \quad P_f(x) + \frac{1}{n}\left\{a_p P_{f+6}(x) + b_p P_{f+4}(x) + c_p P_{f+2}(x) + d_p P_f(x)\right\} \\ &\quad +O(n^{-2}), \end{aligned} \tag{1.1}$$

where

$$\begin{aligned} a_p &= \frac{1}{12}(p^3 + 3p^2 - 12 - 200p^{-1}),\ b_p = \frac{1}{8}(-2p^3 - 5p^2 + 7p - 12 - 420p^{-1}), \\ c_p &= \frac{1}{4}(p^3 + 2p^2 - p - 2 - 216p^{-1}),\ d_p = \frac{1}{24}(-2p^3 - 3p^2 + p + 436p^{-1}). \end{aligned}$$

The first purpose of the paper is to give an analysis of the LRT and John's test for sphericity of large-dimensional observations. It has been well-known that classical multivariate procedures are in general challenged by large-dimensional data. Consequently, modifications and corrections are needed. The reader can find in Johnstone and Titterington (2009) for a review and in Ledoit and Wolf (2002), Srivastava (2005), Srivastava and Fujikoshi (2006), Schott (2007), Bai et al. (2009), Chen et al. (2010), Chen and Qin (2010), Bai et al. (2012) for some representative developments on the topic.

Let us first examine the behavior of the LRT and John's test for large-dimensional data. Empirical sizes of these tests from a Monte-Carlo study are reported in Table 1 where normal vectors with mean zero and identity covariance matrix are used. The sample size is set to $n = 64$ while dimension p increases from 4 to 60 (we have also run other experiments with larger sample sizes n but

TABLE 1

Empirical sizes of LRT and John's test at 5% *significance level based on* 10000 *independent replications using normal vectors* $\mathcal{N}(0, I_p)$ *for* $n = 64$ *and different values of* p.

(p, n)	(4,64)	(8,64)	(16,64)	(32,64)	(48,64)	(56,64)	(60,64)
LRT	0.0483	0.0480	0.0489	0.0520	0.0814	0.2922	0.6673
Nagao's test	0.0468	0.0492	0.0474	0.0487	0.0494	0.0507	0.0535

observations are very similar). The nominal level equals $\alpha = 0.05$ and empirical sizes are computed from 10000 independent replications for each pair of (p, n).

We observe that the Box-Bartlett correction of LRT remains efficient for dimensions $p \leq n/2$ approximatively and, for larger p the test quickly becomes inconsistent with empirical sizes much larger than the nominal level (%5). In contrast, John's test is remarkably robust against the dimension inflation.

In this paper, we propose new corrections to both the LRT and John's test which are consistent within the large-dimensional context. These corrections are valid for general observations with finite fourth moments (not necessarily normal). In the case of John's test, our findings give a new confirmation of its robustness, a result previously found in Ledoit and Wolf (2002) but under the normal assumption.

From the technical point of view, our approach differs from Ledoit and Wolf (2002) and follows the one devised in Bai et al. (2009) and Bai et al. (2012). The central tool is a CLT for linear spectral statistics of sample covariance matrices established in Bai and Silverstein (2004) and later refined in Pan and Zhou (2008). The paper also contains an original contribution on this CLT reported in the Appendix: new formula for the limiting parameters in the CLT are established. Since such CLT's are increasingly important in large-dimensional statistics, we believe that these new formula will be of independent interests for applications other than those considered in the paper.

The remaining of the paper is organized as follows. Large-dimensional corrections to the LRT and John's test are introduced in Section 2. Next, Section 3 reports a detailed Monte-Carlo study to analyze finite-sample sizes and powers of these two corrected large-dimensional tests under both normal and non-normal distributed data. Technical proofs and calculations are relegated to Section 4.

The last section contains some concluding remarks.

2. Large-dimensional corrections

From now on, we assume that the observations $Y_1, \ldots, Y_n$ have the representation $Y_j = \Sigma_p^{1/2} X_j$ where the $p \times n$ table $\{X_1, \ldots, X_n\} = \{x_{ij}\}_{1 \le i \le p, 1 \le j \le n}$ are made with an array of i.i.d. standardized random variables (mean 0 and variance 1). This setting is motivated by the random matrix theory and it is generic enough for a precise analysis of the sphericity test. Furthermore, under the null hypothesis $H_0 : \Sigma_p = \sigma^2 I_p$ (σ^2 is unspecified), we notice that both the LRT and John's test are invariant tests. In particular, the test statistics are independent from the scale parameter σ^2 under the null. Therefore, we can assume w.l.o.g. $\sigma^2 = 1$ when dealing with the null distributions of these test statistics. This will be assumed in all the sections.

Throughout the paper we will use an indicator κ set to 2 when $\{x_{ij}\}$ are real and to 1 when they are complex. Define the kurtosis coefficient $\beta = E|x_{ij}|^4 - 1 - \kappa$ for both cases and note that for normal variables, $\beta = 0$.

2.1. The corrected likelihood ratio test

For the LRT, let $\mathcal{L}_n = -2n^{-1} \log L_n$ for $n \ge 1$. Our first main result is the following.

Theorem 2.1. *Assume $\{x_{ij}\}$ are iid, satisfying $Ex_{ij} = 0, E|x_{ij}|^2 = 1, E|x_{ij}|^4 < \infty$. Then under H_0 and when $\frac{p}{n} = y_n \to y \in (0, 1)$,*

$$\mathcal{L}_n + (p-n) \cdot \log(1 - \frac{p}{n}) - p \Longrightarrow N\{-\frac{\kappa - 1}{2} \log(1-y) + \frac{1}{2}\beta y, -\kappa \log(1-y) - \kappa y\} .$$

The test based on this asymptotic normal distribution will be hereafter referred as *the corrected likelihood-ratio test* (CLRT). One may observe that the limiting distribution of the test crucially depends on the limiting dimension-to-sample ratio y through the factor $-\log(1-y)$. In particular, the asymptotic variance will blow up quickly when y approaches 1. In this case, it is expected that the power of the test will seriously break down. Monte-Carlo experiments in Section 3 will provide more details on this behavior.

The proof of Theorem 2.1 is based on the following lemma. In all the following, F^y denotes the Marčenko-Pastur distribution of index y (> 0) which is introduced and discussed in the Appendix.

Lemma 2.1. *Let* $\{\ell_i\}_{1\le i\le p}$ *be the eigenvalues of the sample covariance matrix* $S_n = n^{-1}\sum_i Y_iY_i^*$. *Then under* H_0 *and the conditions of Theorem 2.1, we have*

$$\begin{pmatrix} \sum_{i=1}^p \log \ell_i - pF^{y_n}(\log x) \\ \sum_{i=1}^p \ell_i - pF^{y_n}(x) \end{pmatrix} \Longrightarrow N(\mu_1, V_1),$$

with

$$\mu_1 = \begin{pmatrix} \frac{\kappa-1}{2}\log(1-y) - \frac{1}{2}\beta y \\ 0 \end{pmatrix} ,$$

and

$$V_1 = \begin{pmatrix} -\kappa\log(1-y) + \beta y & (\beta+\kappa)y \\ (\beta+\kappa)y & (\beta+\kappa)y \end{pmatrix} .$$

The proof of this lemma is postponed to Section 4.

Proof. (of Theorem 2.1). Let $A_n \equiv \sum_{i=1}^p \log \ell_i - pF^{y_n}(\log x)$ and $B_n \equiv \sum_{i=1}^p \ell_i - pF^{y_n}(x)$. By Lemma 2.1,

$$\begin{pmatrix} A_n \\ B_n \end{pmatrix} \Longrightarrow N(\mu_1, V_1).$$

Consequently, $-A_n + B_n$ is asymptotically normal with mean $-\frac{\kappa-1}{2}\log(1-y) + \frac{1}{2}\beta y$ and variance

$$V_1(1,1) + V_1(2,2) - 2V_1(1,2) = -\kappa\log(1-y) - \kappa y.$$

Besides,

$$\begin{aligned}\mathcal{L}_n &= -\Sigma_{i=1}^{p}\log\ell_i + p\log(\frac{1}{p}\Sigma_{i=1}^{p}\ell_i)\\ &= -(A_n + pF^{y_n}(\log x)) + p\log(\frac{1}{p}(B_n+p))\\ &= -A_n - pF^{y_n}(\log x) + p\log(1+\frac{B_n}{p}).\end{aligned}$$

Since $B_n \Rightarrow N(0, y(\beta+\kappa))$, $B_n = O_p(1)$ and $\log(1+B_n/p) = B_n/p + O_p(1/p^2)$. Therefore,

$$\mathcal{L}_n = -A_n - pF^{y_n}(\log x) + B_n + O_p(\frac{1}{p})\ .$$

The conclusion follows with the following well-known integrals w.r.t. the Marčenko-Pastur distribution F^y $(y<1)$,

$$F^y(\log x) = \frac{y-1}{y}\log(1-y) - 1\ , \quad F^y(x) = 1.$$

The proof of Theorem 2.1 is complete. □

2.2. The corrected John's test

Earlier than the asymptotic expansion (1.1) given in Nagao (1973), John (1971) proved that when the observations are normal, the sphericity test based on T_2 is a locally most powerful invariant test. He also established in John (1972) that under these conditions, the limiting distribution of T_2 under H_0 is χ_f^2 with degree of freedom $f = \frac{1}{2}p(p+1) - 1$, or equivalently,

$$nU - p \Longrightarrow \frac{2}{p}\chi_f^2 - p\ ,$$

where for convenience in this section, we have let $U = 2(np)^{-1}T_2$. Clearly, this limit has been established for $n \to \infty$ and a fixed dimension p. However, if we now let $p \to \infty$ in the right-hand side of the above result, it is not hard to see that $\frac{2}{p}\chi_f^2 - p$ will tend to the normal distribution $N(1,4)$. It then seems "natural" to conjecture that when both p and n grow to infinity in some "proper" way, it may happen that

$$nU - p \Longrightarrow N(1,4)\ . \tag{2.1}$$

This is indeed the main result of Ledoit and Wolf (2002) where this asymptotic distribution was established assuming that data are normal-distributed and p and n grow to infinity in a proportional way (i.e. $p/n \to y > 0$).

In this section, we provide a more general result using our own approach. In particular, the distribution of the observation is arbitrary provide a finite fourth moment exists.

Theorem 2.2. *Assume* $\{x_{ij}\}$ *are iid, satisfying* $Ex_{ij} = 0, E|x_{ij}|^2 = 1, E|x_{ij}|^4 < \infty$*, and let* $U = 2(np)^{-1}T_2$*. Then under* H_0 *and when* $p \to \infty, n \to \infty, \frac{p}{n} = y_n \to y \in (0, \infty)$*,*

$$nU - p - \kappa - \beta \Longrightarrow N(-1, 2\kappa) \ .$$

A striking fact in this theorem is that contrary to the limiting distribution for the corrected LRT, the limiting distribution of John's test is *independent* of the dimension to sample ratio $y = \lim p/n$. In this sense, Theorem 2.2 gives a theoretic explanation to the widely observed robustness of John's test against the dimension inflation. Moreover, John's test is also valid for the p larger (or much larger) than n case in contrast to the LRT where this ratio should be kept smaller than 1 to avoid null eigenvalues.

It is also worth noticing that for real normal data, we have $\kappa = 2$ and $\beta = 0$ so that the theorem above reduces to $nU - p \Rightarrow N(1, 4)$. This is exactly the result discussed above from Ledoit and Wolf (2002). Besides, if the data has a non-normal distribution but having the same first four moments as the normal distribution, we have again $nU - p \Rightarrow N(1, 4)$.

The test based on the asymptotic normal distribution given in Theorem 2.2 will be hereafter referred as the *corrected John's test.* Note that in practice for the application of the test with real data, we may estimate the parameter β using the fourth-order sample moment:

$$\widehat{\beta} = \frac{1}{np} \sum_{i=1}^{p} \sum_{j=1}^{n} |Y_j(i)|^4 - \kappa - 1 \ .$$

The proof of Theorem 2.2 is based on the following lemma.

Lemma 2.2. *Let* $\{\ell_i\}_{1\le i\le p}$ *be the eigenvalues of the sample covariance matrix* $S_n = n^{-1}\sum_i Y_i Y_i^*$. *Then under* H_0 *and the conditions of Theorem 2.2, we have*

$$\begin{pmatrix} \sum_{i=1}^p \ell_i^2 - p(1+y_n) \\ \sum_{i=1}^p \ell_i - p \end{pmatrix} \Longrightarrow N(\mu_2, V_2),$$

with

$$\mu_2 = \begin{pmatrix} (\kappa-1+\beta)y \\ 0 \end{pmatrix} ,$$

and

$$V_2 = \begin{pmatrix} 2\kappa y^2 + 4(\kappa+\beta)(y+2y^2+y^3) & 2(\kappa+\beta)(y+y^2) \\ 2(\kappa+\beta)(y+y^2) & (\kappa+\beta)y \end{pmatrix} .$$

The proof of this Lemma is postponed to Section 4.

Proof. (of Theorem 2.2). The result of Lemma 2.2 can be rewritten as:

$$n\begin{pmatrix} p^{-1}\sum_{i=1}^p \ell_i^2 - 1 - \frac{p}{n} - \frac{\kappa+\beta-1}{n} \\ p^{-1}\sum_{i=1}^p \ell_i - 1 \end{pmatrix} \Longrightarrow N\Bigg(\begin{pmatrix} 0 \\ 0 \end{pmatrix}, \frac{1}{y^2}\cdot V_2\Bigg).$$

Define the function $f(x,y) = \frac{x}{y^2} - 1$, then $U = f(p^{-1}\Sigma_{i=1}^p \ell_i^2,\ p^{-1}\Sigma_{i=1}^p \ell_i)$. We have

$$\begin{aligned} &\frac{\partial f}{\partial x}(1+\frac{p}{n}+\frac{\kappa+\beta-1}{n},1) = 1 , \\ &\frac{\partial f}{\partial y}(1+\frac{p}{n}+\frac{\kappa+\beta-1}{n},1) = -2(1+\frac{p}{n}+\frac{\kappa+\beta-1}{n}) , \\ &f(1+\frac{p}{n}+\frac{\kappa+\beta-1}{n},1) = \frac{p}{n}+\frac{\kappa+\beta-1}{n} . \end{aligned}$$

By the delta method,

$$n\Big(U - f(1+\frac{p}{n}+\frac{\kappa+\beta-1}{n},1)\Big) \Longrightarrow N(0, \lim C),$$

where

$$C = \begin{pmatrix} \frac{\partial f}{\partial x}(1+\frac{p}{n}+\frac{\kappa+\beta-1}{n},1) \\ \frac{\partial f}{\partial y}(1+\frac{p}{n}+\frac{\kappa+\beta-1}{n},1) \end{pmatrix}^T \cdot \Big(\frac{1}{y^2}V_2\Big) \cdot \begin{pmatrix} \frac{\partial f}{\partial x}(1+\frac{p}{n}+\frac{\kappa+\beta-1}{n},1) \\ \frac{\partial f}{\partial y}(1+\frac{p}{n}+\frac{\kappa+\beta-1}{n},1) \end{pmatrix}$$

$$\longrightarrow 2\kappa \ .$$

Therefore,

$$n(U - \frac{p}{n} - \frac{\kappa + \beta - 1}{n}) \Longrightarrow N(0, 2\kappa) \ ,$$

that is,

$$nU - p - \kappa - \beta \Longrightarrow N(-1, 2\kappa) \ .$$

The proof of Theorem 2.2 is complete. □

3. Monte Carlo study

Monte Carlo simulations are conducted to find empirical sizes and powers of the corrected likelihood ratio test (CLRT) and the corrected John's test (CJ). In particular, here we want to examine the following questions: how robust are the tests against non-normal distributed data and what is the range of the dimension to sample ratio p/n where the tests are applicable?

More recently, Srivastava (2005) puts forward a statistic T_1 to test the sphericity. Further in the paper Srivastava et al. (2011), they exam the performance of T_1 under non-normality. As the condition $\gamma = 3 + O(p^{-\epsilon})$ in Srivastava et al. (2011) reduces to $\gamma \to 3$ when the dimension p tends to infinity, which is a restriction of the fourth moment matching the normal distribution. Besides, their result is the same as : $nU - p - 1 \Longrightarrow N(0, 4)$ in the normal case (a small perturbation is allowed), which is equivalent to LW's result. So we do not compare their result with our ones in the following.

Chen et al. (2010) propose another test statistic to test the sphericity following the idea of Ledoit and Wolf (2002) but meanwhile they consider some higher order terms in order to estimate $tr(\Sigma)$ and $tr(\Sigma^2)$.

Let

$$Y_{1,n} = \frac{1}{n}\sum_{i=1}^{n} X_i^{'}X_i \ , \qquad Y_{2,n} = \frac{1}{P_n^2}\sum_{i\neq j}(X_i^{'}X_j)^2 \ ,$$
$$Y_{3,n} = \frac{1}{P_n^2}\sum_{i\neq j} X_i^{'}X_j \ , \qquad Y_{4,n} = \frac{1}{P_n^3}\sum_{i,j,k}^{*} X_i^{'}X_jX_j^{'}X_k \ ,$$
$$Y_{5,n} = \frac{1}{P_n^4}\sum_{i,j,k,l}^{*} X_i^{'}X_jX_k^{'}X_l \ ,$$

and define

$$T_{1,n} = Y_{1,n} - Y_{3,n} \ ,$$
$$T_{2,n} = Y_{2,n} - 2Y_{4,n} + Y_{5,n} \ ,$$

then under the sphericity hypothesis, they prove that

$$pn\Big(\frac{T_{2,n}}{T_{1,n}^2}\Big) - n \Longrightarrow N(0,4) \tag{3.1}$$

without distributional assumption for the data, which is exactly the model we are considering here and we refer to this test as C test.

For comparison, we show both the performance of the LW test using the asymptotic $N(1,4)$ distribution in (2.1) (Notice however this is the CJ test under normal distributions) and the C test using the asymptotic $N(0,4)$ distribution in (3.1). The nominal test level is set to be $\alpha = 0.05$, and for each pair of (p, n), we run 10000 independent replications.

We consider two scenarios with respect to the random vectors Y_i :

(a) Y_i is p-dimensional real random vector from the normal population $N(0, I_p)$. In this case, $\kappa = 2$ and $\beta = 0$.

(b) Y_i consists of iid real random variables with distribution $\Gamma(4,2) - 2$ so that y_{ij} satisfies $Ey_{ij} = 0, Ey_{ij}^4 = 4.5$. In this case, $\kappa = 2$ and $\beta = 1.5$.

Table 2 reports the sizes of the four statistics in these two scenarios for different values of (p, n). We see that when $\{y_{ij}\}$ are normal, both LW (=CJ), CLRT and C have similar empirical sizes tending to the nominal level 0.05 as

Table 2

Empirical sizes of LW, CJ, CLRT and C test at 5% *significance level based on 10000 independent applications with real* $N(0,1)$ *random variables and with real Gamma(4,2)-2 random variables.*

(p,n)	$N(0,1)$			Gamma(4,2)-2			
	LW/CJ	CLRT	C	LW	CLRT	CJ	C
(4,64)	0.0424	0.0472	0.494	0.1102	0.1566	0.1577	0.1469
(8,64)	0.0546	0.0577	0.0594	0.131	0.1136	0.1234	0.1139
(16,64)	0.0538	0.056	0.0589	0.1373	0.0836	0.1002	0.0896
(32,64)	0.0533	0.0553	0.0586	0.134	0.0635	0.0804	0.0709
(48,64)	0.0527	0.0507	0.0559	0.1343	0.0608	0.0781	0.0665
(56,64)	0.0542	0.0533	0.0603	0.1301	0.0603	0.0761	0.0657
(60,64)	0.0558	0.0589	0.0576	0.1306	0.059	0.0729	0.0681
(8,128)	0.0547	0.055	0.0584	0.134	0.1165	0.1198	0.1125
(16,128)	0.0523	0.0519	0.054	0.1343	0.0862	0.0954	0.0891
(32,128)	0.0548	0.0532	0.0566	0.1347	0.0691	0.0819	0.0768
(64,128)	0.0555	0.0556	0.0575	0.1254	0.0572	0.0705	0.0655
(96,128)	0.0532	0.0554	0.0557	0.1226	0.0521	0.0626	0.0608
(112,128)	0.0522	0.0518	0.0553	0.123	0.0532	0.0632	0.0618
(120,128)	0.0545	0.0554	0.0557	0.1221	0.0535	0.0625	0.0623
(16,256)	0.0577	0.0587	0.0598	0.1324	0.0869	0.0919	0.0912
(32,256)	0.0579	0.0552	0.0583	0.1272	0.0668	0.071	0.0691
(64,256)	0.0512	0.0553	0.0523	0.1256	0.058	0.0618	0.0596
(128,256)	0.0523	0.0517	0.0533	0.1216	0.0565	0.0555	0.0557
(192,256)	0.0508	0.0522	0.0523	0.12	0.0549	0.0524	0.0508
(224,256)	0.0494	0.052	0.051	0.1169	0.053	0.0537	0.0532
(240,256)	0.0496	0.052	0.051	0.1166	0.0532	0.0517	0.0525
(32,512)	0.0543	0.0555	0.0558	0.1234	0.0663	0.069	0.0676
(64,512)	0.0534	0.052	0.0542	0.1212	0.0567	0.0594	0.0589
(128,512)	0.053	0.0502	0.0527	0.1147	0.0544	0.0529	0.0529
(256,512)	0.0522	0.0522	0.0523	0.1152	0.0558	0.0558	0.0544
(384,512)	0.0484	0.0493	0.05	0.1159	0.0528	0.0524	0.0503
(448,512)	0.0514	0.0497	0.0514	0.1143	0.0511	0.0511	0.0514
(480,512)	0.0529	0.0501	0.054	0.1116	0.0502	0.0549	0.0521

either p or n increases. But when $\{y_{ij}\}$ are Gamma-distributed, the sizes of the statistic LW are higher than 0.1 no matter how large the values of p and n are while the sizes of CLRT and CJ all converge to the nominal level 0.05 as either p or n gets larger. This empirically confirms that normal assumptions are needed for the result of Ledoit and Wolf (2002) while our corrected statistics CLRT and CJ (also the C test) have no such distributional restriction.

Besides, for small values of n, such as $n = 64$ or $n = 128$, the sizes of CLRT are a little bit closer to 0.05 than those of CJ, the main reason is that in the proof of Theorem 2.2, we take the limit of C (the asymptotic variance there), in which we regard the $O(1/yn^2)$ term as $o(1)$. So for any $y \in (0, \infty)$, the larger the value of n, the better the performance is. Another way of saying is that CJ is more sensitive to small values of n. Consider the same procedure we do in the proof of Theorem 2.1 for CLRT, where we regard $\log(1 + B_n/p)$ as B_n/p plus a small term $O(1/p^2)$, and this small term is eventually discarded for large values of p.

As for empirical powers, we consider two alternatives :

(1) Σ_p is diagonal with half of its diagonal elements 0.5 and half 1. We denote its power by Power 1 ;

(2) Σ_p is diagonal with 1/4 of the elements equal 0.5 and 3/4 equal 1. We denote its power by Power 2 .

Table 3 reports the powers of LW(=CJ), CLRT and C when $\{y_{ij}\}$ are distributed as $N(0, 1)$, and of CJ, CLRT and C when $\{y_{ij}\}$ are distributed as Gamma(4,2)-2, for the situation when n equals 64 or 128, with varying values of p and under the above mentioned two alternatives. For $n = 256$ and p varying from 16 to 240, all the tests have powers around 1 under both alternatives so that these values are omitted. And in order to find the trend of these powers, we also present the results when $n = 128$ in Figure 1 and Figure 2.

The behavior of Power 1 and Power 2 in each figure related to the three statistics are similar, except that Power 1 is much higher compared with Power 2 for a given dimension design (p, n) and any given test for the reason that the

Table 3

Empirical powers of LW, CJ, CLRT and C test at 5% *significance level based on 10000 independent applications with real* $N(0,1)$ *random variables and with real Gamma(4,2)-2 random variables under two alternatives Power 1 and 2 (see the text for details).*

$N(0,1)$

(p,n)	Power 1			Power 2		
	LW/CJ	CLRT	C	LW/CJ	CLRT	C
(4,64)	0.7172	0.7411	0.711	0.3814	0.5255	0.3805
(8,64)	0.8123	0.8227	0.8042	0.4303	0.5844	0.4278
(16,64)	0.8645	0.8547	0.8523	0.4597	0.5908	0.4592
(32,64)	0.8991	0.811	0.89	0.475	0.5084	0.4715
(48,64)	0.9092	0.6751	0.8992	0.4825	0.3821	0.4834
(56,64)	0.9111	0.5477	0.9021	0.4908	0.2919	0.4842
(60,64)	0.9112	0.4345	0.9014	0.4908	0.2312	0.4841
(8,128)	0.9972	0.9971	0.997	0.9071	0.9639	0.9032
(16,128)	0.9997	0.9998	0.9998	0.9427	0.9863	0.9392
(32,128)	1	1	1	0.9556	0.9894	0.9525
(64,128)	1	1	1	0.9629	0.9773	0.9609
(96,128)	1	0.9989	1	0.9644	0.9143	0.9621
(112,128)	1	0.9863	0.9999	0.9649	0.8065	0.9598
(120,128)	1	0.9522	1	0.9672	0.695	0.9644

$Gamma(4,2)-2$

(p,n)	Power 1			Power 2		
	CJ	CLRT	C	CJ	CLRT	C
(4,64)	0.5996	0.6335	0.6164	0.3478	0.4708	0.3649
(8,64)	0.714	0.7379	0.7226	0.3996	0.5189	0.4085
(16,64)	0.7912	0.7829	0.7921	0.4396	0.5439	0.4406
(32,64)	0.8559	0.7712	0.8541	0.4815	0.4913	0.4663
(48,64)	0.8837	0.6587	0.8777	0.4904	0.3755	0.4788
(56,64)	0.8838	0.5371	0.88	0.4899	0.2941	0.4791
(60,64)	0.8856	0.4339	0.884	0.4876	0.2368	0.4763
(8,128)	0.9789	0.9824	0.9796	0.8239	0.9021	0.8301
(16,128)	0.9971	0.9976	0.9973	0.8901	0.9537	0.8919
(32,128)	0.9996	0.9998	0.9997	0.9286	0.9745	0.9275
(64,128)	1	0.9997	1	0.9495	0.9672	0.9488
(96,128)	0.9999	0.9977	1	0.9586	0.9037	0.9552
(112,128)	1	0.983	1	0.9606	0.8026	0.9558
(120,128)	1	0.9482	1	0.9617	0.6824	0.9567

first alternative differs more from the null than the second one. The powers of LW (in the normal case), CJ (in the Gamma case) and C are all monotonically increasing in p for a given fixed value of n. But for CLRT, when n is fixed, the powers first increase in p and then become decreasing when p is getting close to n. This can be explained by the fact that when p is close to n, some of the eigenvalues of S_n are getting close to zero, causing the CLRT nearly degenerate and losing power.

Besides, from these figures, we can find that in the normal case the trend of C's power is very much alike of those of LW while in the Gamma case it is similar with those of CJ under both these two alternatives. And in most of the cases, the power of C test is slightly lower than LW (in the normal case) and CJ (in the Gamma case).



Figure 1: Empirical powers of LW/CJ, CLRT and C test at 5% significance level based on 10000 independent applications with real N(0,1) random variables for fixed $n = 128$ under two alternatives Power 1 and 2 (see the text for details).

Lastly, we examine the performance of CJ and C when p is larger than n and empirical sizes and powers are presented in Table 4. We choose the variables to be distributed as Gamma(4,2)-2 since CJ reduces to LW in the normal case, and Ledoit and Wolf (2002) has already reported the performance of LW when



Figure 2: Empirical powers of CJ, CLRT and C test at 5% significance level based on 10000 independent applications with real Gamma(4,2)-2 random variables for fixed $n = 128$ under two alternatives Power 1 and 2 (see the text for details).

p is larger than n. From the table, we see that when p is larger than n, the size of CJ is still correct and it converges to the nominal level 0.05 as the dimension p increases and the same phenomenon exists for C test.

When we evaluate the power, the same two alternatives Power 1 and Power 2 as above are considered. The sample size is fixed to $n = 64$ and the ratio p/n varies from 1 to 20. From the table, we see that Power 1 are much higher than Power 2 for the same reason that the first alternative is easier to be distinguished from H_0. Besides, the powers under both alternatives all increase monotonically (although quite slowly in the case of Power 2) as the dimension p increases (however the power has a small drop in the last case of $(p, n) = (1280, 64)$ and we are suspecting that this is due to some numerical inaccuracy of eigenvalue computation for large p). Besides, the powers of CJ are always slightly higher than those of C in this "large p small n" setting.

An interesting and somewhat surprising phenomenon here is that although asymptotic distribution for the CLRT and CJ are derived under the "Marcenko-Pasture scheme" (i.e $p/n \to c > 0$), the simulation results confirm that even for "large p small n" situation such as $(p, n) = (1280, 64)$, the tests still behave well, which turns out to be free from the assumption constraint "$p/n \to c''$.

Summarizing all these findings from this Monte-Carlo study, the overall figure is the following: when the ratio p/n is much lower than 1 (say smaller than 1/2), it is preferable to employ CLRT (than CJ, LW or C); while this ratio is higher, CJ (or LW for normal data) becomes more powerful (slightly more powerful than C).

Table 4

Empirical sizes and powers (Power 1 and 2) of CJ test and C test at 5% *significance level based on 10000 independent applications with real Gamma(4,2)-2 random variables when* $p \geq n$.

(p, n)	CJ			C		
	Size	Power 1	Power 2	Size	Power 1	Power 2
(64,64)	0.0732	0.8883	0.4869	0.069	0.8838	0.4736
(320,64)	0.0631	0.9208	0.501	0.0587	0.9129	0.4891
(640,64)	0.0587	0.9296	0.5141	0.0552	0.9186	0.493
(960,64)	0.0579	0.9315	0.5143	0.0544	0.9202	0.5006
(1280,64)	0.0599	0.9284	0.5088	0.0544	0.9178	0.4916

4. Additional proofs

4.1. Proof of Lemma 2.1

Let for $x > 0$, $f(x) = \log x$ and $g(x) = x$. Define A_n and B_n by the decompositions

$$\begin{aligned}\sum_{i=1}^{p} \log \ell_i &= p \int f(x) d(F_n(x) - F^{y_n}(x)) + pF^{y_n}(f) = A_n + pF^{y_n}(f) \ , \\ \sum_{i=1}^{p} \ell_i &= p \int g(x) d(F_n(x) - F^{y_n}(x)) + pF^{y_n}(g) = B_n + pF^{y_n}(g) \ .\end{aligned}$$

Applying Theorem 2.1 given in the Appendix to the pair (f, g), we have

$$\begin{pmatrix} A_n \\ B_n \end{pmatrix} \Longrightarrow N\left(\begin{pmatrix} EX_f \\ EX_g \end{pmatrix} \begin{pmatrix} \mathrm{Cov}(X_f, X_f) & \mathrm{Cov}(X_f, X_g) \\ \mathrm{Cov}(X_g, X_f) & \mathrm{Cov}(X_g, X_g) \end{pmatrix} \right).$$

It remains to evaluate the limiting parameters and this results from the following calculations:

$$\begin{aligned} I_1(f,r) &= \frac{1}{2}\log\left(1-h^2/r^2\right)\ , & (4.1)\\ I_1(g,r) &= 0\ , & (4.2)\\ I_2(f) &= -\frac{1}{2}h^2\ , & (4.3)\\ I_2(g) &= 0\ , & (4.4)\\ J_1(f,g,r) &= \frac{h^2}{r^2}\ , & (4.5)\\ J_1(f,f,r) &= -\frac{1}{r}\log(1-h^2/r)\ , & (4.6)\\ J_1(g,g,r) &= \frac{h^2}{r^2}\ , & (4.7)\\ J_2(f,g) &= h^2\ , & (4.8)\\ J_2(f,f) &= h^2\ , & (4.9)\\ J_2(g,g) &= h^2\ . & (4.10) \end{aligned}$$

We now detail these calculations to complete the proof. They are all based on the formula given in Proposition A.1 in the Appendix and repeated use of the residue theorem.

Proof of (4.1): We have

$$\begin{aligned} I_1(f,r) &= \frac{1}{2\pi i}\oint_{|\xi|=1} f(|1+h\xi|^2)\Big[\frac{\xi}{\xi^2-r^{-2}}-\frac{1}{\xi}\Big]d\xi\\ &= \frac{1}{2\pi i}\oint_{|\xi|=1} \log(|1+h\xi|^2)\Big[\frac{\xi}{\xi^2-r^{-2}}-\frac{1}{\xi}\Big]d\xi\\ &= \frac{1}{2\pi i}\oint_{|\xi|=1} (\frac{1}{2}\log((1+h\xi)^2)+\frac{1}{2}\log((1+h\xi^{-1})^2)\Big[\frac{\xi}{\xi^2-r^{-2}}-\frac{1}{\xi}\Big]d\xi\\ &= \frac{1}{2\pi i}\Big[\oint_{|\xi|=1}\log(1+h\xi)\frac{\xi}{\xi^2-r^{-2}}d\xi-\oint_{|\xi|=1}\log(1+h\xi)\frac{1}{\xi}d\xi\\ &\quad+\oint_{|\xi|=1}\log(1+h\xi^{-1})\frac{\xi}{\xi^2-r^{-2}}d\xi-\oint_{|\xi|=1}\log(1+h\xi^{-1})\frac{1}{\xi}d\xi\Big] \end{aligned}$$

For the first integral, note that as $r>1$, the poles are $\pm\frac{1}{r}$ and we have by the residue theorem,

$$\begin{aligned}
&\frac{1}{2\pi i}\oint_{|\xi|=1}\log(1+h\xi)\frac{\xi}{\xi^2-r^{-2}}d\xi \\
=\ &\left.\frac{\log(1+h\xi)\cdot\xi}{\xi-r^{-1}}\right|_{\xi=-r^{-1}}+\left.\frac{\log(1+h\xi)\cdot\xi}{\xi+r^{-1}}\right|_{\xi=r^{-1}} \\
=\ &\frac{1}{2}\log(1-\frac{h^2}{r^2})\ .
\end{aligned}$$

For the second integral,

$$\frac{1}{2\pi i}\oint_{|\xi|=1}\log(1+h\xi)\frac{1}{\xi}d\xi=\log(1+h\xi)\big|_{\xi=0}=0\ .$$

The third one is

$$\begin{aligned}
&\frac{1}{2\pi i}\oint_{|\xi|=1}\log(1+h\xi^{-1})\frac{\xi}{\xi^2-r^{-2}}d\xi \\
&=-\frac{1}{2\pi i}\oint_{|z|=1}\log(1+hz)\frac{z^{-1}}{z^{-2}-r^{-2}}\cdot\frac{-1}{z^2}dz \\
&=\frac{1}{2\pi i}\oint_{|z|=1}\frac{\log(1+hz)r^2}{z(z+r)(z-r)}dz=\left.\frac{\log(1+hz)r^2}{(z+r)(z-r)}\right|_{z=0}=0\ ,
\end{aligned}$$

where the first equality results from the change of variable $z=\frac{1}{\xi}$, and the third equality holds because $r>1$, so the only pole is $z=0$.

The fourth one equals

$$\begin{aligned}
&\frac{1}{2\pi i}\oint_{|\xi|=1}\log(1+h\xi^{-1})\frac{1}{\xi}d\xi=-\frac{1}{2\pi i}\oint_{|z|=1}\log(1+hz)\frac{-z}{z^2}dz \\
&=\log(1+hz)\big|_{z=0}=0\ .
\end{aligned}$$

Collecting the four integrals leads to the desired formula for $I_1(f,r)$.

Proof of (4.2): We have

$$
\begin{aligned}
I_1(g,r) &= \frac{1}{2\pi i}\oint_{|\xi|=1} g(|1+h\xi|^2)[\frac{\xi}{\xi^2-r^{-2}}-\frac{1}{\xi}]d\xi \\
&= \frac{1}{2\pi i}\oint_{|\xi|=1} |1+h\xi|^2\Big[\frac{\xi}{\xi^2-r^{-2}}-\frac{1}{\xi}\Big]d\xi \\
&= \frac{1}{2\pi i}\Big[\oint_{|\xi|=1}(1+h\xi)\frac{\xi}{\xi^2-r^{-2}}d\xi-\oint_{|\xi|=1}\frac{1+h\xi}{\xi}d\xi \\
&\quad +\oint_{|\xi|=1}(1+h\overline{\xi})\frac{\xi}{\xi^2-r^{-2}}d\xi-\oint_{|\xi|=1}\frac{1+h\overline{\xi}}{\xi}d\xi\Big]\ .
\end{aligned}
$$

These four integrals are calculated as follows:

$$
\begin{aligned}
&\frac{1}{2\pi i}\oint_{|\xi|=1}(1+h\xi)\frac{\xi}{\xi^2-r^{-2}}d\xi=\frac{(1+h\xi)\xi}{\xi+r^{-1}}\bigg|_{\xi=\frac{1}{r}}+\frac{(1+h\xi)\xi}{\xi-r^{-1}}\bigg|_{\xi=-\frac{1}{r}}=1\ ; \\
&\frac{1}{2\pi i}\oint_{|\xi|=1}\frac{1+h\xi}{\xi}d\xi=1+h\xi\Big|_{\xi=0}=1\ ; \\
&\frac{1}{2\pi i}\oint_{|\xi|=1}\frac{\xi(1+h\overline{\xi})}{\xi^2-r^{-2}}d\xi=\frac{1}{2\pi i}\oint_{|\xi|=1}\frac{\xi+h}{\xi^2-r^{-2}}d\xi \\
&=\frac{\xi+h}{\xi-r^{-1}}\Big|_{\xi=-r^{-1}}+\frac{\xi+h}{\xi+r^{-1}}\Big|_{\xi=r^{-1}}=1\ ; \\
&\frac{1}{2\pi i}\oint_{|\xi|=1}\frac{1+h\overline{\xi}}{\xi}d\xi=\frac{1}{2\pi i}\oint_{|\xi|=1}\frac{\xi+h}{\xi^2}d\xi=\frac{\partial}{\partial\xi}(\xi+h)\Big|_{\xi=0}=1\ .
\end{aligned}
$$

Collecting the four terms leads to $I_1(g,r)=0$.

Proof of (4.3):

$$
\begin{aligned}
I_2(f) &= \frac{1}{2\pi i}\oint_{|\xi|=1}\log(|1+h\xi|^2)\frac{1}{\xi^3}d\xi \\
&= \frac{1}{2\pi i}\left[\oint_{|\xi|=1}\frac{\log(1+h\xi)}{\xi^3}d\xi+\oint_{|\xi|=1}\frac{\log(1+h\xi^{-1})}{\xi^3}d\xi\right]\ .
\end{aligned}
$$

We have

$$
\begin{aligned}
&\frac{1}{2\pi i}\oint_{|\xi|=1}\frac{\log(1+h\xi)}{\xi^3}d\xi=\frac{1}{2}\frac{\partial^2}{\partial\xi^2}\log(1+h\xi)\Big|_{\xi=0}=-\frac{1}{2}h^2\ ; \\
&\frac{1}{2\pi i}\oint_{|\xi|=1}\frac{\log(1+h\xi^{-1})}{\xi^3}d\xi=-\frac{1}{2\pi i}\oint_{|z|=1}\frac{\log(1+hz)}{\frac{1}{z^3}}\cdot\frac{-1}{z^2}dz=0\ .
\end{aligned}
$$

Combining the two leads to $I_2(f) = -\frac{1}{2}h^2$.

Proof of (4.4):

$$I_2(g) = \frac{1}{2\pi i}\oint_{|\xi|=1} \frac{(1+h\overline{\xi})(1+h\xi)}{\xi^3}d\xi = \frac{1}{2\pi i}\oint_{|\xi|=1} \frac{\xi + h\xi^2 + h + h^2\xi}{\xi^4}d\xi = 0 \ .$$

Proof of (4.5):

$$J_1(f,g,r) \quad = \quad \frac{1}{2\pi i}\oint_{|\xi_2|=1} |1+h\xi_2|^2 \cdot \frac{1}{2\pi i}\oint_{|\xi_1|=1} \frac{\log(|1+h\xi_1|^2)}{(\xi_1 - r\xi_2)^2}d\xi_1 d\xi_2 \ .$$

We have,

$$\begin{aligned} & \frac{1}{2\pi i}\oint_{|\xi_1|=1} \frac{\log(|1+h\xi_1|^2)}{(\xi_1 - r\xi_2)^2}d\xi_1 \\ = \quad & \frac{1}{2\pi i}\oint_{|\xi_1|=1} \frac{\log(1+h\xi_1)}{(\xi_1 - r\xi_2)^2}d\xi_1 + \frac{1}{2\pi i}\oint_{|\xi_1|=1} \frac{\log(1+h\xi_1^{-1})}{(\xi_1 - r\xi_2)^2}d\xi_1 \ . \end{aligned}$$

The first term

$$\frac{1}{2\pi i}\oint_{|\xi_1|=1} \frac{\log(1+h\xi_1)}{(\xi_1 - r\xi_2)^2}d\xi_1 = 0,$$

because for fixed $|\xi_2| = 1$, $|r\xi_2| = |r| > 1$, so $r\xi_2$ is not a pole.

The second term is

$$\begin{aligned} & \frac{1}{2\pi i}\oint_{|\xi_1|=1} \frac{\log(1+h\xi_1^{-1})}{(\xi_1 - r\xi_2)^2}d\xi_1 = -\frac{1}{2\pi i}\oint_{|z|=1} \frac{\log(1+hz)}{(\frac{1}{z} - r\xi_2)^2} \cdot \frac{-1}{z^2}dz \\ & = \frac{1}{2\pi i} \cdot \frac{1}{(r\xi_2)^2}\oint_{|z|=1} \frac{\log(1+hz)}{(z - \frac{1}{r\xi_2})^2}dz = \frac{1}{(r\xi_2)^2} \cdot \frac{\partial}{\partial z}\log(1+hz)\Big|_{z=\frac{1}{r\xi_2}} \\ & = \frac{h}{r\xi_2(r\xi_2 + h)} \ , \end{aligned}$$

where the first equality results from the change of variable $z = \frac{1}{\xi_1}$, and the third equality holds because for fixed $|\xi_2| = 1$, $|\frac{1}{r\xi_2}| = \frac{1}{|r|} < 1$, so $\frac{1}{r\xi_2}$ is a pole of second order.

Therefore,

$$\begin{aligned}
&J_1(f,g,r)\\
=\ &\frac{h}{2\pi i r^2}\oint_{|\xi_2|=1}\frac{(1+h\xi_2)(1+h\overline{\xi_2})}{\xi_2(\xi_2+\frac{h}{r})}d\xi_2\\
=\ &\frac{h}{2\pi i r^2}\oint_{|\xi_2|=1}\frac{\xi_2+h\xi_2^2+h+h^2\xi_2}{\xi_2^2(\xi_2+\frac{h}{r})}d\xi_2\\
=\ &\frac{h}{2\pi i r^2}\left[\oint_{|\xi_2|=1}\frac{1+h^2}{\xi_2(\xi_2+\frac{h}{r})}d\xi_2+\oint_{|\xi_2|=1}\frac{h}{\xi_2+\frac{h}{r}}d\xi_2+\oint_{|\xi_2|=1}\frac{h}{\xi_2^2(\xi_2+\frac{h}{r})}d\xi_2\right].
\end{aligned}$$

Finally we find $J_1(f,g,r)=\frac{h^2}{r^2}$ since

$$\begin{aligned}
&\frac{h}{2\pi i r^2}\oint_{|\xi_2|=1}\frac{1+h^2}{\xi_2(\xi_2+\frac{h}{r})}d\xi_2=0\ ,\quad \frac{h}{2\pi i r^2}\oint_{|\xi_2|=1}\frac{h}{\xi_2+\frac{h}{r}}d\xi_2=\frac{h^2}{r^2}\ ,\\
&\frac{h}{2\pi i r^2}\oint_{|\xi_2|=1}\frac{h}{\xi_2^2(\xi_2+\frac{h}{r})}d\xi_2=0\ .
\end{aligned}$$

Proof of (4.6):

$$\begin{aligned}
J_1(f,f,r) &= \frac{1}{2\pi i}\oint_{|\xi_2|=1} f(|1+h\xi_2|^2)\cdot\frac{1}{2\pi i}\oint_{|\xi_1|=1}\frac{f(|1+h\xi_1|^2)}{(\xi_1-r\xi_2)^2}d\xi_1 d\xi_2\\
&= \frac{1}{2\pi i}\oint_{|\xi_2|=1} f(|1+h\xi_2|^2)\frac{h}{r\xi_2(r\xi_2+h)}d\xi_2\\
&= \frac{h}{2\pi i r^2}\oint_{|\xi_2|=1}\frac{\log(1+h\xi_2)}{\xi_2(\frac{h}{r}+\xi_2)}d\xi_2+\frac{h}{2\pi i r^2}\oint_{|\xi_2|=1}\frac{\log(1+h\xi_2^{-1})}{\xi_2(\frac{h}{r}+\xi_2)}d\xi_2\ .
\end{aligned}$$

We have

$$\begin{aligned}
&\frac{h}{2\pi i r^2}\oint_{|\xi_2|=1}\frac{\log(1+h\xi_2)}{\xi_2(\frac{h}{r}+\xi_2)}d\xi_2\\
=\ &\frac{h}{r^2}\left[\left.\frac{\log(1+h\xi_2)}{\frac{h}{r}+\xi_2}\right|_{\xi_2=0}+\left.\frac{\log(1+h\xi_2)}{\xi_2}\right|_{\xi_2=-\frac{h}{r}}\right]\\
=\ &-\frac{1}{r}\log(1-\frac{h^2}{r})\ ,
\end{aligned}$$

and

$$\begin{aligned}
&\frac{h}{2\pi i r^2}\oint_{|\xi_2|=1}\frac{\log(1+h\xi_2^{-1})}{\xi_2(\frac{h}{r}+\xi_2)}d\xi_2=\frac{-h}{2\pi i r^2}\oint_{|z|=1}\frac{\log(1+hz)}{\frac{1}{z}(\frac{h}{r}+\frac{1}{z})}\cdot\frac{-1}{z^2}dz\\
&=\frac{1}{2\pi i r}\oint_{|z|=1}\frac{\log(1+hz)}{z+\frac{r}{h}}dz=0\ ,
\end{aligned}$$

where the first equality results from the change of variable $z = \frac{1}{\xi_2}$, and the third equality holds because $|\frac{r}{h}| > 1$, so $\frac{r}{h}$ is not a pole.
Finally, we find $J_1(f, f, r) = -\frac{1}{r}\log(1 - \frac{h^2}{r})$.

Proof of (4.7):

$$J_1(g,g,r) = \frac{1}{2\pi i}\oint_{|\xi_2|=1} |1+h\xi_2|^2 \cdot \frac{1}{2\pi i}\oint_{|\xi_1|=1} \frac{|1+h\xi_1|^2}{(\xi_1 - r\xi_2)^2} d\xi_1 d\xi_2 \ .$$

We have

$$\begin{aligned}
&\frac{1}{2\pi i}\oint_{|\xi_1|=1} \frac{|1+h\xi_1|^2}{(\xi_1 - r\xi_2)^2} d\xi_1 = \frac{1}{2\pi i}\oint_{|\xi_1|=1} \frac{\xi_1 + h\xi_1^2 + h + h^2\xi_1}{\xi_1(\xi_1 - r\xi_2)^2} d\xi_1 \\
&= \frac{1}{2\pi i}\Bigg[\oint_{|\xi_1|=1} \frac{1+h^2}{(\xi_1 - r\xi_2)^2} d\xi_1 + \oint_{|\xi_1|=1} \frac{h\xi_1}{(\xi_1 - r\xi_2)^2} d\xi_1 \\
&+ \oint_{|\xi_1|=1} \frac{h}{\xi_1(\xi_1 - r\xi_2)^2} d\xi_1\Bigg] \\
&= \frac{h}{r^2\xi_2^2} \ ,
\end{aligned}$$

since

$$\begin{aligned}
&\frac{1}{2\pi i}\oint_{|\xi_1|=1} \frac{1+h^2}{(\xi_1 - r\xi_2)^2} d\xi_1 = 0 \ , \quad \frac{1}{2\pi i}\oint_{|\xi_1|=1} \frac{h\xi_1}{(\xi_1 - r\xi_2)^2} d\xi_1 = 0 \ , \\
&\frac{1}{2\pi i}\oint_{|\xi_1|=1} \frac{h}{\xi_1(\xi_1 - r\xi_2)^2} d\xi_1 = \left.\frac{h}{(\xi_1 - r\xi_2)^2}\right|_{\xi_1=0} = \frac{h}{r^2\xi_2^2} \ ,
\end{aligned}$$

where the equality above holds because for fixed $|\xi_2| = 1$, $|r\xi_2| = |r| > 1$, so $r\xi_2$ is not a pole. Therefore,

$$\begin{aligned}
J_1(g,g,r) &= \frac{h}{2\pi i r^2}\oint_{|\xi_2|=1} \frac{\xi_2 + h\xi_2^2 + h + h^2\xi_2}{\xi_2^3} d\xi_2 \\
&= \frac{h}{2\pi i r^2}\left[\oint_{|\xi_2|=1} \frac{1+h^2}{\xi_2^2} d\xi_2 + \oint_{|\xi_2|=1} \frac{h}{\xi_2} d\xi_2 + \oint_{|\xi_2|=1} \frac{h}{\xi_2^3} d\xi_2\right] , \\
&= \frac{h^2}{r^2} \ .
\end{aligned}$$

Proof of (4.8) (4.9) (4.10): We have

$$\begin{aligned}&\frac{1}{2\pi i}\oint_{|\xi_1|=1}\frac{f(|1+h\xi_1|^2)}{\xi_1^2}d\xi_1=\frac{1}{2\pi i}\oint_{|\xi_1|=1}\frac{\log(|1+h\xi_1|^2)}{\xi_1^2}d\xi_1\\&=\frac{1}{2\pi i}\oint_{|\xi_1|=1}\frac{\log(1+h\xi_1)+\log(1+h\xi_1^{-1})}{\xi_1^2}d\xi_1\ =h\ ,\end{aligned}$$

since

$$\begin{aligned}&\frac{1}{2\pi i}\oint_{|\xi_1|=1}\frac{\log(1+h\xi_1)}{\xi_1^2}d\xi_1=\frac{\partial}{\partial\xi_1}\Big(\log(1+h\xi_1)\Big)\Big|_{\xi_1=0}=h\ ,\\&\frac{1}{2\pi i}\oint_{|\xi_1|=1}\frac{\log(1+h\xi_1^{-1})}{\xi_1^2}d\xi_1=-\frac{1}{2\pi i}\oint_{|z|=1}\frac{\log(1+hz)}{\frac{1}{z^2}}\cdot(-\frac{1}{z^2}dz)\\&=\frac{1}{2\pi i}\oint_{|z|=1}\log(1+hz)dz=0\ .\end{aligned}$$

Similarly,

$$\frac{1}{2\pi i}\oint_{|\xi_2|=1}\frac{g(|1+h\xi_2|^2)}{\xi_2^2}d\xi_2=\frac{1}{2\pi i}\oint_{|\xi_2|=1}\frac{\xi_2+h\xi_2^2+h+h^2\xi_2}{\xi_2^3}d\xi_2=h.$$

Therefore,

$$\begin{aligned}J_2(f,g)&=\frac{1}{2\pi i}\oint_{|\xi_1|=1}\frac{f(|1+h\xi_1|^2)}{\xi_1^2}d\xi_1\cdot\frac{1}{2\pi i}\oint_{|\xi_2|=1}\frac{g(|1+h\xi_2|^2)}{\xi_2^2}d\xi_2=h^2\ ,\\J_2(f,f)&=\frac{1}{2\pi i}\oint_{|\xi_1|=1}\frac{f(|1+h\xi_1|^2)}{\xi_1^2}d\xi_1\cdot\frac{1}{2\pi i}\oint_{|\xi_2|=1}\frac{f(|1+h\xi_2|^2)}{\xi_2^2}d\xi_2=h^2\ ,\\J_2(g,g)&=\frac{1}{2\pi i}\oint_{|\xi_1|=1}\frac{g(|1+h\xi_1|^2)}{\xi_1^2}d\xi_1\cdot\frac{1}{2\pi i}\oint_{|\xi_2|=1}\frac{g(|1+h\xi_2|^2)}{\xi_2^2}d\xi_2=h^2\ .\end{aligned}$$

The proof of Lemma 2.1 is complete.

4.2. Proof of Lemma 2.2

Let $f(x)=x^2$ and $g(x)=x$. Define C_n and B_n by the decompositions

$$\begin{aligned}\sum_{i=1}^p\ell_i^2&=p\int f(x)d(F_n(x)-F^{y_n}(x))+pF^{y_n}(f)=C_n+pF^{y_n}(f)\ ,\\\sum_{i=1}^p\ell_i&=p\int g(x)d(F_n(x)-F^{y_n}(x))+pF^{y_n}(g)=B_n+pF^{y_n}(g)\ .\end{aligned}$$

Applying Theorem 2.1 given in the Appendix to the pair (f, g), we have

$$\begin{pmatrix} C_n \\ B_n \end{pmatrix} \Longrightarrow N\left(\begin{pmatrix} EX_f \\ EX_g \end{pmatrix} \begin{pmatrix} \mathrm{Cov}(X_f, X_f) & \mathrm{Cov}(X_f, X_g) \\ \mathrm{Cov}(X_g, X_f) & \mathrm{Cov}(X_g, X_g) \end{pmatrix} \right).$$

It remains to evaluate the limiting parameters and this results from the following calculations:

$$I_1(f, r) = \frac{h^2}{r^2}, \tag{4.11}$$

$$I_1(g, r) = 0, \tag{4.12}$$

$$I_2(f) = h^2, \tag{4.13}$$

$$I_2(g) = 0, \tag{4.14}$$

$$J_1(f, g, r) = \frac{2h^2 + 2h^4}{r^2}, \tag{4.15}$$

$$J_1(f, f, r) = \frac{2h^4 + (2h + 2h^3)^2 r}{r^3}, \tag{4.16}$$

$$J_1(g, g, r) = \frac{h^2}{r^2}, \tag{4.17}$$

$$J_2(f, g) = 2h^2 + 2h^4, \tag{4.18}$$

$$J_2(f, f) = (2h + 2h^3)^2, \tag{4.19}$$

$$J_2(g, g) = h^2. \tag{4.20}$$

The results (4.12), (4.14), (4.17) and (4.20) are exactly the same as those found in (4.2), (4.4), (4.7) and (4.10) in the proof of Lemma 2.1. The remaining results are found by similar calculations using Proposition A.1 in the Appendix and their details are omitted (available however in a separated on-line supplementary file).

5. Concluding remarks

Using recent central limit theorems for eigenvalues of large sample covariance matrices, we are able to find new asymptotic distributions for two major procedures to test the sphericity of a large-dimensional distribution. Although the theory is developed under the scheme $p \to \infty$, $n \to \infty$ and $p/n \to y > 0$, our Monte-Carlo study has proved that: on the one hand, both the corrected LRT and the corrected John's test are already very efficient for middle dimension such as $(p, n) = (96, 128)$ both in size and power, see Table 2 and Table 3, and on the other hand, the corrected John's test also behaves very well in most of "large p, small n" situation, see Table 4.

Three characteristic features emerge from our findings:

(a) These asymptotic distributions are universal in the sense that they depend on the distribution of the observations only through its first four moments;

(b) The new test procedures improve quickly when either the dimension p or the sample size n gets large. In particular, for a given sample size n, higher dimensions p lead to better performance of these corrected test statistics.

(c) The corrected John's test is particularly robust against the dimension inflation. Our Monte-Carlo study shows that for a small sample size $n = 64$, the test is effective for $0 < p/n \leq 20$.

In a sense, these new procedures have benefited from the "blessings of dimensionality".

References

Anderson, T.W. (1984). *An introduction to Multivariate Statistical Analysis*, 2nd ed. Wiley, New York.

Bai, Z.D. and Silverstein, J.W. (2004). CLT for linear spectral statistics of large-dimensional sample covariance matrices. *Ann. Probab.*, **32**, 553–605.

Bai, Z., Jiang, D., Yao, J. and Zheng, S. (2009). Corrections to LRT on large-dimensional covariance matrix by RMT. *Ann. Statist.*, 37, 3822–3840.

Bai, Z.D. and Silverstein, J.W. (2010). *Spectral Analysis of Large Dimensional Random Matrices* (2nd edition). Springer, 20.

Bai, Z., Jiang, D., Yao, J. and Zheng, S. (2012). Testing linear hypotheses in high-dimensional regressions. *Statistics*, 2012 (On-line First).

Chen, S.X. and Qin, Y.L. (2010). A two-sample test for high-dimensional data with applications to gene-set testing. *Ann. Statist.*, 38, 808–835.

Chen, S.X., Zhang, L.X. and Zhong, P.S. (2010). Tests for high-dimensional covariance matrices. *J. Amer. Statist. Assoc.*, 105, 810–819.

John, S. (1971). Some optimal multivariate tests. *Biometrika,* **58**, 123–127.

John, S. (1972). The distribution of a statistic used for testing sphericity of normal distributions. *Biometrika,* **59**, 169–173.

Jonsson, D. (1982). Some limit theorems for the eigenvalues of a sample covariance matrix. *J. Multivariate Anal.*, **12**(1), 1–38.

Johnstone, I.M. and Titterington, D.M. (2009). Statistical challenges of high-dimensional data. *Philos. Trans. R. Soc. Lond. Ser. A Math. Phys. Eng. Sci.*, **367**, 4237–4253.

Ledoit, O. and Wolf, M. (2002). Some hypothesis tests for the covariance matrix when the dimension is large compared to the sample size. *Ann. Statist.*, **30**, 1081–1102.

Lytova, A. and Pastur, L. (2009). Central limit theorem for linear eigenvalue statistics of the Wigner and the sample covariance random matrices. *Metrika*, **69**, 153–172.

Mauchly, J.W. (1940). Test for sphericity of a normal n-variate distribution. *Ann. Mathe. Statist.*, **11**, 204–209.

Muirhead, R.J. (1982). *Aspects of Multivariate Statistical Theory.* Wiley, New York.

Nagao, H. (1973). On some test criteria for covariance matrix. *Ann. Statist.*, **1**, 700–709.

Pan, G.M. and Zhou, W. (2008). Central limit theorem for signal-to-interference ratio of reduced rank linear receiver. *Ann. Appl. Probab.*, **18**, 1232-1270.

Schott, J.R. (2007). Some high-dimensional tests for a one-way MANOVA. *J. Multivariate Anal.*, 98, 1825–1839.

Srivastava, M.S. (2005). Some tests concerning the covariance matrix in high dimensional data. *J. Japan Statist. Soc.*, 35, 251–272.

Srivastava, M.S. and Fujikoshi, Y. (2006). Multivariate analysis of variance with fewer observations than the dimension. *J. Multiv. Anal.*, 97, 1927–1940.

Srivastava, M.S. Kollo, T. and Rosen, D. (2011). Some tests for the covariance matrix with fewer observations than the dimension under non-normality. *J. Multiv. Anal.*, 102, 1090–1103.

Sugiura, N. (1972). Locally best invariant test for sphericity and the limiting distributions. *Ann. Mathe. Statist.*, **43**, 1312-1316.

Zheng, S. (2012). Central Limit Theorem for Linear Spectral Statistics of Large Dimensional F Matrix. *Ann. Institut Henri Poincaré Probab. Statist.*, **48**, 444–476.

Appendix A: Formula for limiting parameters in the CLT for eigenvalues of a sample covariance matrix with general fourth moments

Given a sample covariance matrix S_n of dimension p with eigenvalues $\lambda_1, \ldots, \lambda_p$, linear spectral statistics of the form $F_n(g) = p^{-1} \sum_{i=1}^{p} g(\lambda_j)$ for suitable functions g are of central importance in multivariate analysis. Such CLT's have been successively developed since the pioneering work of Jonsson (1982), see Bai and Silverstein (2004) and Lytova and Pastur (2009) for a recent account on the subject.

The CLT in Bai and Silverstein (2004) (see also an improved version in Bai and Silverstein (2010)) has been widely used in applications as this CLT also provides, for the first time, explicit formula for the mean and covariance parameters of the normal limiting distribution. In the special case with an array $\{x_{ij}\}$ of independent variables, this CLT assumes the following moment conditions:

(a) For each n, $x_{ij} = x_{ij}^n, i \leqslant p, j \leqslant n$ are independent.

(b) $Ex_{ij} = 0, E|x_{ij}|^2 = 1, \max_{i,j,n} E|x_{ij}|^4 < \infty$.

(c) If $\{x_{ij}\}$'s are real, then $Ex_{ij}^4 = 3$; otherwise (complex variables), $Ex_{ij}^2 = 0$ and $E|x_{ij}|^4 = 2$.

In Condition (c), the fourth moments of the entries are set to the values 3 or 2 matching the normal case. This is indeed a quite demanding and restrictive condition since in the real case for example, it is incredibly hard to find a non-normal distribution with mean 0, variance 1 and fourth moment equaling 3. As a consequence, most of if not all applications published in the literature using this CLT assumes a normal distribution for the observations. Recently, effort have been made in Pan and Zhou (2008), Lytova and Pastur (2009) and Zheng (2012) to overcome these moment restrictions. We present below such a CLT with general forth moments that will be used for the sphericity test.

In all the following, we use an indicator κ set to 2 when $\{x_{ij}\}$ are real and to 1 when they are complex. Define $\beta = E|x_{ij}|^4 - 1 - \kappa$ for both cases and $h = \sqrt{y}$.

For the presentation of the results, let be the sample covariance matrix $S_n = \frac{1}{n}\sum_{i=1}^n X_i X_i^*$ where $X_i = (x_{ki})_{1\le k\le p}$ is the i-th observed vector. It is then well-known that when $p \to \infty$, $n \to \infty$ and $p/n \to y > 0$, the distribution of its eigenvalues converges to a nonrandom distribution, namely the Marčenko-Pastur distribution F^y with support $[a, b] = [(1 \pm \sqrt{y})^2]$ (an additional mass at the origin when $y > 1$). Moreover, the Stieltjes transform $\underline{m}$ of a companion distribution defined by $\underline{F}^y = (1-y)\delta_0 + yF_c$ satisfies an inverse equation for $z \in \mathbb{C}^+$,

$$z = -\frac{1}{\underline{m}} + \frac{y}{1+\underline{m}}. \tag{A.1}$$

The following CLT is a particular instance of Theorem 1.4 in Pan and Zhou (2008).

Theorem A.1. *[Pan and Zhou (2008)] Assume that for each* n*, the variables* $x_{ij} = x_{ij}^n, i \leqslant p, j \leqslant n$ *are independent and identically distributed satisfying* $Ex_{ij} = 0$*,* $E|x_{ij}|^2 = 1$*,* $E|x_{ij}|^4 = \beta + 1 + \kappa < \infty$ *and in case of they are complex,* $Ex_{ij}^2 = 0$*. Assume moreover,*

$$p \to \infty, \quad n \to \infty, \quad p/n \to y > 0\ .$$

Let $f_1, \cdots f_k$ *be functions analytic on an open region containing the support of* F^y*. The random vector* $\{X_n(f_1), \cdots X_n(f_k)\}$ *where*

$$X_n(f) = p\left\{F_n(f) - F^{y_n}(f)\right\}$$

converges weakly to a normal vector $(X_{f_1},\cdots X_{f_k})$ *with mean function and covariance function:*

$$\begin{aligned}\mathbb{E}[X_f] &= (\kappa-1)I_1(f)+\beta I_2(f)\ , && \text{(A.2)}\\ Cov(X_f,X_g) &= \kappa J_1(f,g)+\beta J_2(f,g)\ , && \text{(A.3)}\end{aligned}$$

where

$$\begin{aligned}I_1(f) &= -\frac{1}{2\pi i}\oint \frac{y\left\{\underline{m}/(1+\underline{m})\right\}^3(z)f(z)}{\left[1-y\left\{\underline{m}/(1+\underline{m})\right\}^2\right]^2}dz\ ,\\ I_2(f) &= -\frac{1}{2\pi i}\oint \frac{y\left\{\underline{m}/(1+\underline{m})\right\}^3(z)f(z)}{1-y\left\{\underline{m}/(1+\underline{m})\right\}^2}dz\ ,\\ J_1(f,g) &= -\frac{1}{4\pi^2}\oint\oint \frac{f(z_1)g(z_2)}{(\underline{m}(z_1)-\underline{m}(z_2))^2}\underline{m}'(z_1)\underline{m}'(z_2)dz_1dz_2\ ,\\ J_2(f,g) &= \frac{-y}{4\pi^2}\oint f(z_1)\frac{\partial}{\partial z_1}\left\{\frac{\underline{m}}{1+\underline{m}}(z_1)\right\}dz_1\cdot\oint g(z_2)\frac{\partial}{\partial z_2}\left\{\frac{\underline{m}}{1+\underline{m}}(z_2)\right\}dz_2\ ,\end{aligned}$$

where the integrals are along contours (non overlapping in J_1*) enclosing the support of* F^y*.*

However, concrete applications of this CLT are not easy since the limiting parameters are given through those integrals on contours that are only vaguely defined. The purpose of this appendix is to go a step further by providing alternative formula for these limiting parameters. These new formula, presented in the following Proposition convert all these integral along the unit circle; they are much easier to use for concrete applications, see for instance the proofs of Lemma 2.1 and 2.2 in the paper. Furthermore, these formula will be of independent interest for applications other than those developed in this paper.

Proposition A.1. *The limiting parameters in Theorem A.1 can be expressed* *as following:*

$$
\begin{aligned}
I_1(f) &= \lim_{r\downarrow 1} I_1(f,r)\ , &\text{(A.4)}\\
I_2(f) &= \frac{1}{2\pi i}\oint_{|\xi|=1} f(|1+h\xi|^2)\frac{1}{\xi^3}d\xi\ , &\text{(A.5)}\\
J_1(f,g) &= \lim_{r\downarrow 1} J_1(f,g,r)\ , &\text{(A.6)}\\
J_2(f,g) &= -\frac{1}{4\pi^2}\oint_{|\xi_1|=1}\frac{f(|1+h\xi_1|^2)}{\xi_1^2}d\xi_1\oint_{|\xi_2|=1}\frac{g(|1+h\xi_2|^2)}{\xi_2^2}d\xi_2\ , &\text{(A.7)}
\end{aligned}
$$

with

$$
\begin{aligned}
I_1(f,r) &= \frac{1}{2\pi i}\oint_{|\xi|=1} f(|1+h\xi|^2)\left[\frac{\xi}{\xi^2-r^{-2}}-\frac{1}{\xi}\right]d\xi\ ,\\
J_1(f,g,r) &= -\frac{1}{4\pi^2}\oint_{|\xi_1|=1}\oint_{|\xi_2|=1}\frac{f(|1+h\xi_1|^2)g(|1+h\xi_2|^2)}{(\xi_1-r\xi_2)^2}d\xi_1 d\xi_2\ .
\end{aligned}
$$

Proof. We start with the simplest formula $I_2(f)$ to explain the main argument and indeed, the other formula are obtained similarly. The idea is to introduce the change of the variable $z = 1+hr\xi+hr^{-1}\overline{\xi}+h^2$ with $r>1$ but close to 1 and $|\xi|=1$ (recall $h=\sqrt{y}$). Note that this idea has been employed in Zheng (2012). It can be readily checked that when ξ runs counterclockwisely the unit circle, z will run a contour $\mathcal{C}$ that encloses closely the support interval $[a,b]=[(1\pm h)^2]$ (recall $h=\sqrt{y}$). Moreover, by the Eq. (A.1), we have on $\mathcal{C}$

$$
\underline{m} = -\frac{1}{1+hr\xi},\quad \text{and}\quad dz = h(r-r^{-1}\xi^{-2})d\xi\ .
$$

Applying this variable change to the formula of $I_2(f)$ given in Theorem A.1, we have

$$
\begin{aligned}
I_2(f) &= \lim_{r\downarrow 1}\frac{1}{2\pi i}\oint_{|\xi|=1} f(z)\frac{1}{\xi^3}\frac{r\xi^2-r^{-1}}{r(r^2\xi^2-1)}d\xi\\
&= \frac{1}{2\pi i}\oint_{|\xi|=1} f(|1+h\xi|^2)\frac{1}{\xi^3}d\xi\ .
\end{aligned}
$$

This proves the formula (A.5). For (A.4), we have similarly

$$
\begin{aligned}
I_1(f) &= \lim_{r\downarrow 1}\frac{1}{2\pi i}\oint_{|\xi|=1} f(z)\frac{1}{\xi^3}\frac{r\xi^2-r^{-1}}{r(r^2\xi^2-1)}\frac{1}{1-r^{-2}\xi^{-2}}d\xi\\
&= \lim_{r\downarrow 1}\frac{1}{2\pi i}\oint_{|\xi|=1} f(|1+h\xi|^2)\frac{1}{\xi(\xi^2-r^{-2})}\\
&= \lim_{r\downarrow 1} I_1(f,r)\ .
\end{aligned}
$$

Formula (A.7) for $J_2(f,g)$ is calculated in a same fashion by observing that we have

$$\frac{\partial}{\partial z}\left\{\frac{\underline{m}}{1+\underline{m}}(z)\right\}dz = \frac{\partial}{\partial \xi}\left\{\frac{\underline{m}}{1+\underline{m}}(\xi)\right\}d\xi = \frac{\partial}{\partial \xi}\left\{\frac{1}{-hr\xi}\right\}d\xi = \frac{1}{hr\xi^2}d\xi \ .$$

Finally for (A.6), we use two non-overlapping contours defined by $z_j = 1 + hr_j\xi_j + hr_j^{-1}\overline{\xi}_j + h^2$, $j = 1,2$ where $r_2 > r_1 > 1$. By observing that

$$\underline{m}'(z_j)dz_j = \left(\frac{\partial}{\partial \xi_j}\underline{m}\right)d\xi_j = \frac{hr_j}{(1+hr_j\xi_j)^2}d\xi_j \ ,$$

we find

$$\begin{aligned} J_1(f,g) &= \lim_{\substack{r_2>r_1>1\\ r_2\downarrow 1}} -\frac{1}{4\pi^2}\oint_{|\xi_1|=1}\oint_{|\xi_2|=1}\frac{f(z_1)g(z_2)}{\{\underline{m}(z_1)-\underline{m}(z_2)\}^2} \\ &\quad \cdot\frac{hr_1}{(1+hr_1\xi_1)^2}\cdot\frac{hr_2}{(1+hr_2\xi_2)^2}d\xi_1 d\xi_2 \\ &= \lim_{\substack{r_2>r_1>1,\\ r_2\downarrow 1}} -\frac{1}{4\pi^2}\oint_{|\xi_1|=1}\oint_{|\xi_2|=1}\frac{f(z_1)g(z_2)}{\{r_1\xi_1-r_2\xi_2\}^2}d\xi_1 d\xi_2 \\ &= \lim_{r\downarrow 1} -\frac{1}{4\pi^2}\oint_{|\xi_1|=1}\oint_{|\xi_2|=1}\frac{f(|1+h\xi_1|^2)g(|1+h\xi_2|^2)}{\{\xi_1-r\xi_2\}^2}d\xi_1 d\xi_2 \ . \end{aligned}$$

The proof is complete. □